

Reassessing optimal climate-change mitigation strategies through more explicit consideration of the role of time in impact assessments.

Miko U.F. Kirschbaum

**Landcare Research
Palmerston North
New Zealand**

e-mail: KirschbaumM@LandcareResearch.co.nz

Forest → pasture → forest




CO₂ with full carbon cycle

Nitrous oxide (long-lived gas)

Methane (short-lived gas)

Albedo (surface reflectance)

CO₂ emissions are rising fast




Key points:

CO₂ concentrations are rising rapidly, with annual growth rate now being about 2 ppmv yr⁻¹. The atmospheric increase is driven primarily by the large emissions from the burning of fossil fuels. Observed emissions are at the upper range of IPCC scenarios.

Raupach et al. (2007); Steffen (2009); NOAA (2009); IMF (2009)

Nitrous oxide



Key points:

N₂O concentrations have been going up steadily for at least the past 25 years. While concentrations are low, they are nonetheless important because N₂O has powerful radiative absorptive properties.



Methane

IPCC (2007)

Key points:

Methane concentrations have more than doubled since pre-industrial times, then appeared to have stabilised by about 2000. The most recent measurements suggest, however, that its atmospheric concentration has started to increase again.


NOAA (2009)



Forests absorb more radiation than pastures

Key points:

The Earth receives basically all of its energy as short-wave radiation from the sun which must be balanced by long-wave emission back into space. The proportion of short-wave radiation absorbed has a direct effect on the temperature of the Earth. Grassland reflect more incoming radiation than forests (18% vs 13% are typical values under snow-free conditions). The greater radiation absorption of forests has a direct warming effect.




Betts (2000)


Forests absorb more radiation than pastures, especially with a snow cover

Key points:

The albedo effect is strongest when snow covers the ground. Grassland may then reflect $\frac{3}{4}$ instead of $\frac{1}{4}$ of incoming radiation. The warming effect of that extra radiation absorption by forests can then be substantial.




0.78



0.26

Albedo

Betts (2000)




©IPCC 2007: WG1-AR4

Key points:
 This graph shows the contribution to observed radiative forcing changes by all radiative forcing agents. The radiative forcing agents affected by land-use change are circled.

IPCC (2007)

Deforestation

Dairying



Key points:

The graph shows the contribution to global warming for 1 ha converted from forest to dairying. It converts radiative forcing directly into temperature changes and the lag between radiative forcing and temperature changes is ignored in these calculations.

Albedo changes at the time of land-use change and makes an on-going cooling contribution.

N₂O is added year after year to cumulatively add to its warming contribution.

CH₄ adds to warming for some years, but then an equilibrium is reached, with new added CH₄ balancing the loss of methane emitted in earlier years.

CO₂ makes a one-off large contribution when the original forest is cut. Some of that atmospheric CO₂ is taken up by the oceans to lower the warming effect of the originally emitted CO₂ over later years.

Methane: 240 kgCH₄ ha⁻¹ yr⁻¹
Nitrous oxide: 10 kgN₂O (N) ha⁻¹ yr⁻¹
Forest: 200 tC ha⁻¹
Albedo: 0.13 and 0.18 (forest, pasture)

CH₄ and N₂O emission factors from Saggarr (pers. comm.)

Deforestation

Dairying

Integrals over 100 years

CO₂	27.5 (51%)
CH₄	14.9 (28%)
N₂O	11.2 (21%)
Albedo	-6.8 (-13%)
Total	46.7

Key points:

This shows the integrated effect of warming effect in individual years integrated over 100 years. CO₂ contributes about half of total warming and CH₄ and N₂O about a quarter each. Albedo changes offset the warming effect by about 10-20%.

These relativities are strongly affected by the magnitude of respective emissions of the different greenhouse gases. The numbers here are based on New Zealand numbers for dairying, which have high emissions of methane and nitrous oxide. It is also based on large numbers for carbon loss which are based on the emission when a mature *Pinus radiata* stand is felled.

Methane: 240 kgCH₄ ha⁻¹ yr⁻¹

Nitrous oxide: 10 kgN₂O (N) ha⁻¹ yr⁻¹


Forest: 200 tC ha⁻¹

Albedo: 0.13 and 0.18 (forest, pasture)


CH₄ and N₂O emission
factors from Saggar
(pers. comm.)

Reforestation

Wood for paper use



With wood products



Model assumptions:
33-year rotations, albedo increase over 10 years, no methane or nitrous oxide emissions from forests.
Paper use – immediate loss
Wood products: 50% of biomass, exponential decay with 25-yr time constant

Key points:

When pastures are reforested, the C storage in trees has a cooling effect. That is partly negated as albedo decreases, with a warming effect. When trees are harvested, C storage gains are reversed. Atmospheric CO₂ is even higher (with a net-warming effect) than it would have been without temporary carbon storage in trees because ocean uptake is reduced over the time when atmospheric CO₂ is lowered by C storage in trees. This pattern is much less pronounced when longer-lived wood products are produced and extended the period of time over which C is held out of the atmosphere.

Reforestation

Wood for paper use

Actual With credit

CO₂	-12.8	-12.8 (-38%)
CH₄	-0.1	-11.5 (-34%)
N₂O	0	-9.8 (-29%)
Albedo	5.8	5.8 (17%)
Total	-7.1	-28.4

With wood products

Actual With credit

CO₂	-18.1	-18.1 (-46%)
CH₄	-0.1	-11.5 (-29%)
N₂O	0	-9.8 (-25%)
Albedo	5.8	5.8 (15%)
Total	-12.5	-33.7


Key points:

This shows the integrated cooling effect from reforestation in individual years integrated over 100 years. It is shown both for the actual emissions (assuming no methane or nitrous oxide emissions from forests), or with inclusion of credits for avoided emissions that would have occurred from on-going pastoral activity. Looking at the numbers with credit, and for wood products used for paper, CO₂, CH₄ and N₂O each contribute about the same to overall cooling, and albedo changes offset the benefit by about 15-20%.

When long-lived wood products are produced, the benefit of reforestation increases, but the effect is not huge, increasing the carbon benefit by about 50% (from 12.8 to 18.1), but for all radiative agents combined, it increases the benefit by only 15% (from 28.4 to 33.7)

Reforestation

Bioenergy



Key points:

When bioenergy can be generated from wood products (and thereby replace CO₂ release from burning fossil fuels), it has the same benefits as shown on previous slides, and additionally generates on-going and accumulating benefits through fossil-fuel substitution. If further increases the benefit of reforestation.

The increased benefit is also not large, but can accrue beyond the 100-year analysis horizon.

Actual With credit

CO₂	-27.5	-27.5 (-56%)
CH₄	-0.1	-11.5 (-24%)
N₂O	0	-9.8 (-20%)
Albedo	5.8	5.8 (12%)
Total	-7.1	-43.1

Reforestation (100-yr gains) [nK ha⁻¹ (100 yr)⁻¹]

	CO₂	Total
No harvest	-23.6	-38.5
Short-rotation 15 years (paper use)	-5.0	-21.5
Short-rotation 33 years (paper use)	-12.8	-28.4
Short-rotation (construction wood)	-18.1	-33.7
Short rotation (bioenergy)	-27.5	-43.1


Key points:

This summarises the benefit of different reforestation options by considering only CO₂ effects or all effects, including those from avoided pastoral emissions and avoided fossil-fuel emissions.

Short rotations with wood used for paper has the lowest benefit. It increases with length of the rotation, or when wood is used for construction purposes. Wood used for bioenergy has the greatest benefit because of the on-going and accruing substitution of fossil fuels. Over 100 years, the benefit of a bioenergy plantation is comparable to that of a permanently re-established forest.

Differences between options are quite marked if one considers only CO₂ benefits, but options are more similar if one looks at the total effect from all radiative agents as their effect is similar for all forest-use options.

A closer look at time



Key points:

The different greenhouse gases have quite different atmospheric lifetimes. The atmospheric lifetimes of methane and nitrous oxide are usually described with first-order decay kinetics of 12 and 120 years, respectively, and CO₂ is described with multiple decay constants for different fractions of emitted CO₂. These different turn-over times affect their respective climatic impacts.

Which aspect of climate change impacts us most?

Instantaneous climatic conditions?

- Heat damage
- Severe weather
- Tropical diseases (e.g. malaria)
- Food production

Rate of climate change?

- Ecological mal-adaptation
- Socio-economic institutions

Cumulative climate change?





- Sea level rise

Key points:

These different climatic impacts require very different quantifications. Instantaneous climate impacts are quantified directly as a function of temperature itself. Rate of change impacts are quantified as a function of temperature divided by the time over which a temperature increase has occurred.

Cumulative temperature impacts are quantified through summing the temperature increases over the number of years over which they occur.

A detailed look at impacts (from 1 year's worth of emission)



Key points:

The top graphs give the instantaneous and rate of change impact due to a single one-off emission of a unit of greenhouse gas. It shows that N₂O and CO₂ emitted in 2000 still have a substantial warming impact in 2100 whereas all methane emitted in 2000 will have been oxidised well before 2100.

Key points:

The bottom graphs show that in terms of cumulative temperature impacts, a unit of gas emitted of all three gases will add to the cumulative impact experienced in 2100. In the case of CO₂ and N₂O, the cumulative impact will further continue to increase beyond 2100.

Traditional greenhouse warming potentials are calculated with an approach similar to that used to calculate cumulative temperature impacts.

Greenhouse Warming Potentials or Climate Change Impact Potentials (CCIP) (relative to CO₂, 100 year horizon)

Current

CH₄ = 25

N₂O = 298

CCIP

CH₄ ≈ 9

N₂O ≈ 298


Key points:

If one

- defines the mitigation goal to minimise the most severe climatic impacts expected to be experienced by the end of this century; and
- gives each of the three kinds of the climatic impacts (defined above) equal weight, then
- that would not make much difference for the quantification of N₂O relative to CO₂; but
- it would greatly reduce the calculated (impact-weighted) warming potential of methane from 25 to about 9.

This would have significant implications for the relative importance attached to the emission of different greenhouse gases and the urgency in controlling their respective emissions. It would mean that short-term emission controls should focus primarily on the long-lived gases N₂O and CO₂, whereas emission control of methane can be given somewhat lower priority in the near term.

An example: The effect of short-term plantations





10-year plantations on pasture, cease CH₄ and N₂O emissions for 10 years; wood growth for 10 years, then harvested with immediate release of CO₂.

Key points:

This assesses the mitigation effect of planting a forest in 2050, cutting it after 10 years and thereafter immediately releasing the temporarily stored carbon. Atmospheric CO₂ is decreased while the forest is storing carbon, but after re-release of the carbon in the forest, atmospheric CO₂ is higher than it would have been without temporary storage. Hence, atmospheric CO₂ and temperature in 2100 will be higher than without the 10-year plantation. Methane radiative forcing decreases while the land is under forest, but the gains are lost over the following decades. By 2100, there is only a very small benefit remaining. The N₂O concentration falls while land is under forest, and the gain is maintained to 2100 and beyond.

An example: The effect of short-term plantations

Analysed by effect on ameliorating most severe impacts to 2100.



Key points:

It shows that a short-term plantation would marginally **worsen** instantaneous and rate of change impacts related to CO₂, and more so for plantations established later in the century, but it ameliorates cumulative temperature impacts.

For methane, plantations established before about 2050, have almost no effect on instantaneous and rate of change impacts but improve cumulative temperature impacts. Only plantations established later can also improve instantaneous and rate of change impacts.

For nitrous oxide, plantations established at any time affect significant amelioration of all kinds of impacts, and for cumulative temperature impacts, the benefit diminishes almost linearly with any delay in establishing those plantations.

Climate change impacts (the key questions)

Instantaneous climatic conditions?

Rate of climate change?

Cumulative climate change?

Give equal weight? Or is one more important than others?

Minimise the most severe impacts? To 2100?

Impacts weighted (e.g. $I_{\text{targ}} = \sum I^s$, with $s > 1$)?

Apply discount factors? Is the future less important than the present?

Traditional GWPs assume the answers without asking the questions

Conclusions

- **CO₂, CH₄, N₂O are of comparable significance under LUC**
- **Albedo changes are opposite in their effect to that of greenhouse gases by (≈10-20%) and more in snow-covered areas**
- **Impact assessment should quantify impacts in detail, esp. cumulative vs instantaneous**
- **The importance of methane is over-rated.**

References:

- Betts R.A. (2000). Offset of the potential carbon sink from boreal forestation by decreases in surface albedo. *Nature* **408**: 187–190.
- International Monetary Fund (2009) World economic outlook. October 2009.
<http://www.imf.org/external/pubs/ft/weo/2009/02/index.htm>
- IPCC (2007) Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (eds: Solomon S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M.M.B., Miller, H.L. Jr and Chen, Z.) Cambridge University Press, Cambridge, UK and New York, NY, USA. 996 pp.
- Kirschbaum, M.U.F. (2003). Can trees buy time? An assessment of the role of vegetation sinks as part of the global carbon cycle. *Climatic Change* **58**: 47-71.
- Kirschbaum, M.U.F. (2003). To sink or burn? A discussion of the potential contributions of forests to greenhouse gas balances through storing carbon or providing biofuels. *Biomass and Bioenergy* **24**: 297-310.
- Kirschbaum, M.U.F. (2006). Temporary carbon sequestration cannot prevent climate change. *Mitigation and Adaptation Strategies for Global Change* **11**: 1151-1164.
- Kirschbaum, M.U.F., Sagar, S., Tate, K.R., Giltrap, D.L., Ausseil, A.G.E., Greenhalgh, S., Whitehead, D. (2010). Assessing the climate change implications of shifting land use between forest and permanent pasture (In preparation).
- NOAA (2009). <http://climateprogress.org/2009/04/25/noaa-methane-levels-2008/>
- Raupach, M.R., Marland, G., Ciais, P., Le Quéré, C., Canadell, J.G., Klepper, G. and Field, C.B. (2007) Global and regional drivers of accelerating CO₂ emissions. *Proceedings of the National Academy of Sciences (USA)* **104**: 10288–10293.
- Steffen, W. (2009). Climate Change 2009. Faster Change & More Serious Risks. Australian Department of Climate Change, Canberra.